

# Supernova Bound on keV-mass Sterile Neutrinos

Shun Zhou

Max-Planck-Institut für Physik, München

Based on *G.G. Raffelt & S.Z., Phys.Rev.D 83, 093014 (2011)*



HAvSE 2011

Hamburg Neutrinos from Supernova Explosions



# Outline

- 1. Motivation: WDM, Pulsar Kicks**
- 2. Sterile Neutrinos in SN Cores**
- 3. Energy losses and SN bounds**



# Robust evidence for Dark Matter



## Cold or Warm DM ?



## Abundance of Cosmic Substructure



## WDM candidate: keV-mass sterile neutrinos

In the early Universe: *production via neutrino oscillations*

$$\begin{aligned} |\nu_\alpha\rangle &= +\cos\mathcal{G}|\nu_1\rangle + \sin\mathcal{G}|\nu_2\rangle \\ |\nu_s\rangle &= -\sin\mathcal{G}|\nu_1\rangle + \cos\mathcal{G}|\nu_2\rangle \end{aligned}$$

Described by two parameters:  $m_s$  &  $\mathcal{G}$

Boltzmann equation: *distribution function of sterile neutrinos*

$$\frac{\partial}{\partial t} f_s(p, t) - H p \frac{\partial}{\partial p} f_s(p, t) \approx \Gamma(\nu_\alpha \rightarrow \nu_s; p, t) [f_\alpha(p, t) - f_s(p, t)]$$

Matter effects: *primordial lepton number asymmetry*

$$\sin^2 2\theta_m \approx \frac{(\Delta m^2/2p)^2 \sin^2 2\theta}{(\Delta m^2/2p)^2 \sin^2 2\theta + (\Delta m^2/2p \cos 2\theta - V_m - V_T)^2}$$


**Non-resonant case:** Dodelson, Widrow, 94'

**Resonant case:** Shi, Fuller, 99'

$$\Omega_s h^2 \approx 0.1 \left( \frac{\sin^2 \mathcal{G}}{3 \times 10^{-9}} \right) \left( \frac{m_s}{3 \text{ keV}} \right)^{1.8}$$




$$\Omega_s h^2 \approx 0.1 \left( \frac{m_s}{1 \text{ keV}} \right) \left( \frac{L}{3 \times 10^{-3}} \right)$$

# Pulsar velocities: asymmetric emission of sterile neutrinos



Neutrino oscillations in magnetized media



Kusenko, 09'



Hobbs et al., 05'



A statistical study of 233 pulsar proper motions

# X-ray observations: to discover keV-mass sterile neutrinos



# Various constraints on keV-mass sterile neutrinos

Kusenko, 09'





# Standard energy-loss arguments

## Neutrino Cooling

$$\mathcal{E}_\nu = 3.0 \times 10^{33} \text{ erg cm}^{-3} \text{ s}^{-1}$$

If new weakly interacting particles can be produced in the supernova core, they will steal energies from neutrino bursts, which reduces the duration of neutrino signals.



Volume Emission of New Particles

$$\mathcal{E}_{\text{new}} < 3.0 \times 10^{33} \text{ erg cm}^{-3} \text{ s}^{-1}$$

G.G. Raffelt, Phys. Rept. 198 (1990) 1

Proto-Neutron Star

$$\rho \approx \rho_{\text{nuc}} = 3 \times 10^{14} \text{ g cm}^{-3}$$

$$T \approx 30 \text{ MeV}$$

Axions, Majorons, *Sterile Neutrinos*, ...



# Sterile Neutrinos in SN Cores

## Production of sterile neutrinos:

Low matter density: neutrino flavor oscillations with matter effects

High matter density: production & absorption via scattering processes

## Occupation-number formalism:

Sigl, Raffelt, 93'

$$\rho_{ij} = \langle b_i^+ b_j \rangle$$

$$\bar{\rho}_{ij} = \langle d_j^+ d_i \rangle$$

Diagonal terms: just the usual occupation numbers

Non-diagonal terms: encode the phase information

## Equations of motion:

$$\dot{\rho}_{\mathbf{p}} = i[\rho_{\mathbf{p}}, \Omega_{\mathbf{p}}] + \sum_{i=1}^n \left[ \left( I_i - \frac{1}{2} \{I_i, \rho_{\mathbf{p}}\} \right) \mathcal{P}_{\mathbf{p}}^i - \frac{1}{2} \{I_i, \rho_{\mathbf{p}}\} \mathcal{A}_{\mathbf{p}}^i \right] \quad \text{Neutral-current interaction}$$

$$+ \frac{1}{2} \sum_a \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \left[ \mathcal{W}_{\mathbf{p}'\mathbf{p}}^a \left( G^a \rho_{\mathbf{p}'} G^a (1 - \rho_{\mathbf{p}}) + \text{h.c.} \right) - \mathcal{W}_{\mathbf{p}\mathbf{p}'}^a \left( \rho_{\mathbf{p}} G^a (1 - \rho_{\mathbf{p}'}) G^a + \text{h.c.} \right) \right]$$

# Sterile Neutrinos in SN Cores

## Two-flavor mixing case


$$\rho_{\mathbf{p}} = \frac{1}{2} (n_{\mathbf{p}} + \mathbf{P}_{\mathbf{p}} \cdot \boldsymbol{\tau})$$

$$\Omega_{\mathbf{p}} = \frac{1}{2} (E_{\mathbf{p}} + \mathbf{B}_{\mathbf{p}} \cdot \boldsymbol{\tau})$$

$$\dot{\rho}_{\mathbf{p}} = i[\rho_{\mathbf{p}}, \Omega_{\mathbf{p}}] \longrightarrow \dot{\mathbf{P}}_{\mathbf{p}} = \mathbf{B}_{\mathbf{p}} \times \mathbf{P}_{\mathbf{p}}$$

Flavor polarization vectors rotate around magnetic fields

$$\mathbf{B}_{\mathbf{p}} = \left( \frac{\Delta m^2}{2E} \sin 2\vartheta, \quad 0, \quad \frac{\Delta m^2}{2E} \cos 2\vartheta - V_{\text{eff}} \right)$$



## Matter Effects

Wolfenstein, 78'; Mikheyev, Smirnov, 85

$$\sin^2 2\vartheta_{\nu, \bar{\nu}} = \frac{\sin^2 2\vartheta}{\sin^2 2\vartheta + (\cos 2\vartheta \mp (\pm)E / E_r)^2}$$

where the resonant energy is

$$E_r = \frac{\Delta m^2}{2|V_{\text{eff}}|}$$

maximal mixing if  $E \sim E_r \cos 2\vartheta$

# Sterile Neutrinos in SN Cores

## Weak-damping limit


Oscillation length

$$\lambda_{\text{osc}} = \frac{4\pi E}{\Delta\tilde{m}^2} < 0.7 \text{ cm} \left( \frac{E}{30 \text{ MeV}} \right) \left( \frac{10^{-4}}{\sin 2\mathcal{G}} \right) \left( \frac{10 \text{ keV}}{m_s} \right)^2$$



Mean free path

$$\lambda_{\text{mfp}} = \frac{1}{N_B \sigma_{\nu N}} \approx 10^3 \text{ cm} \left( \frac{30 \text{ MeV}}{E} \right)^2 \left( \frac{10^{14} \text{ g cm}^{-3}}{\rho} \right)$$

Neutrinos oscillate many times before a subsequent collision with nucleons



$$\lambda_{\text{osc}} \ll \lambda_{\text{mfp}}$$



# Sterile Neutrinos in SN Cores

## In the weak-damping limit

$$\tilde{\rho}_{\mathbf{p}} = \frac{1}{2} \left[ n_{\mathbf{p}} + (\mathbf{P}_{\mathbf{p}} \cdot \hat{\mathbf{B}}_{\mathbf{p}}) (\hat{\mathbf{B}}_{\mathbf{p}} \cdot \boldsymbol{\tau}) \right]$$

averaged over a period of oscillation

Two independent parameters:  
occupation numbers  $f_{\mathbf{p}}^{\alpha}$  &  $f_{\mathbf{p}}^s$

$$\tilde{\rho}_{\mathbf{p}} = \begin{pmatrix} f_{\mathbf{p}}^{\alpha} & 0 \\ 0 & f_{\mathbf{p}}^s \end{pmatrix} + \frac{1}{2} (f_{\mathbf{p}}^{\alpha} - f_{\mathbf{p}}^s) \boldsymbol{\tau}_{\mathbf{p}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

## Simplified equations of motion:

$$\dot{f}_{\mathbf{p}}^s = \frac{1}{4} s_{\mathbf{p}}^2 \left\{ \left[ (1 - f_{\mathbf{p}}^s) \mathcal{P}_{\mathbf{p}}^{\alpha} - f_{\mathbf{p}}^s \mathbf{a}_{\mathbf{p}}^{\alpha} \right] + \sum_a (g_a^{\alpha})^2 \int \frac{d^3 \mathbf{p}'}{(2\pi)^3} \left[ \mathbf{w}_{\mathbf{p}'\mathbf{p}}^a f_{\mathbf{p}'}^{\alpha} (1 - f_{\mathbf{p}}^s) - \mathbf{w}_{\mathbf{p}\mathbf{p}'}^a f_{\mathbf{p}}^s (1 - f_{\mathbf{p}'}^{\alpha}) \right] \right\}$$

further simplification if sterile neutrinos escape from the SN core

$$\dot{f}_{\mathbf{p}}^s = \frac{1}{4} s_{\mathbf{p}}^2 \left[ \mathcal{P}_{\mathbf{p}}^{\alpha} + \sum_a (g_a^{\alpha})^2 \int \frac{d^3 \mathbf{p}'}{(2\pi)^3} \mathbf{w}_{\mathbf{p}'\mathbf{p}}^a f_{\mathbf{p}'}^{\alpha} \right]$$

set  $f_{\mathbf{p}}^s = 0$

Lepton-number-loss rate  $\dot{\mathcal{N}}_{\text{L}} = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \dot{f}_{\mathbf{p}}^s$

Energy-loss rate  $\dot{\mathcal{E}}_s = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} E f_{\mathbf{p}}^s$

# Sterile Neutrinos in SN Cores

## Neutrino matter potentials

$$V_{\nu_e} = \sqrt{2}G_F N_B \left[ Y_e - \frac{1}{2}Y_n + 2Y_{\nu_e} + Y_{\nu_\mu} + Y_{\nu_\tau} \right]$$

$$V_{\nu_\mu} = \sqrt{2}G_F N_B \left[ -\frac{1}{2}Y_n + Y_{\nu_e} + 2Y_{\nu_\mu} + Y_{\nu_\tau} \right]$$

$$V_{\nu_\tau} = \sqrt{2}G_F N_B \left[ -\frac{1}{2}Y_n + Y_{\nu_e} + Y_{\nu_\mu} + 2Y_{\nu_\tau} \right]$$

### Remarks:

1. degenerate electron neutrinos; the equation of state involved; charged current interactions; so we consider tau neutrinos for simplicity;
2. we assume the SN core to be homogeneous and isotropic.

## Tau-sterile neutrino mixing

$$V_{\nu_\tau} = -\frac{G_F}{\sqrt{2}} N_B (1 - Y_e - 2Y_{\nu_e} - 4Y_{\nu_\tau}) < 0$$

### Initial conditions:

$$Y_e = 0.3, Y_{\nu_e} = 0.07, Y_{\nu_\mu} = Y_{\nu_\tau} = 0$$

$$\sin^2 2\mathcal{G}_{\nu, \bar{\nu}} = \frac{\sin^2 2\mathcal{G}}{\sin^2 2\mathcal{G} + (\cos 2\mathcal{G} \pm E/E_r)^2}$$

1. the MSW resonance occurs in the antineutrino channel;
2. asymmetry between tau neutrinos and antineutrinos.

# Energy-loss Rates and SN Bounds

Simple bounds in the 'vacuum limit':  $E_r \gg E$

$$\sin^2 2\mathcal{G}_{\nu, \bar{\nu}} = \frac{\sin^2 2\mathcal{G}}{\sin^2 2\mathcal{G} + (\cos 2\mathcal{G} \pm E/E_r)^2}$$



$$\mathcal{G}_\nu \approx \mathcal{G}_{\bar{\nu}} \approx \mathcal{G}$$

## Energy-loss rates

$$\mathcal{E}_s = \int_0^\infty \frac{E^2}{2\pi^2} \frac{E}{\exp(E/T)+1} \left( \frac{1}{4} \sin^2 2\mathcal{G} \right) \frac{N_B G_F^2 E^2}{\pi} dE = 4N_B G_F^2 T^6 \mathcal{G}^2$$

$\nu + \bar{\nu}$ 
 $\nu - N$

## Supernova Bound

$$\mathcal{E}_s = 4N_B G_F^2 T^6 \mathcal{G}^2 < \mathcal{E}_\nu = 3.0 \times 10^{33} \text{ erg cm}^{-3} \text{ s}^{-1} \implies \mathcal{G}^2 \leq 10^{-8}$$

$$\rho \approx \rho_{\text{nuc}} = 3 \times 10^{14} \text{ g cm}^{-3}$$

$$T \approx 30 \text{ MeV}$$

Such a simple bound is valid and mass-independent only in the 'vacuum limit'.

# Energy-loss Rates and SN Bounds

## Conditions for the vacuum limit

$$E_r = 3.25 \text{ MeV} \left( \frac{m_s}{10 \text{ keV}} \right)^2 \left( \frac{10^{14} \text{ g cm}^{-3}}{\rho} \right) |Y_0 - Y_{\nu_\tau}|^{-1}$$


$$Y_0 = (1 - Y_e - 2Y_{\nu_e})/4 = 0.14$$

$$m_s > 100 \text{ keV}$$

$$Y_{\nu_\tau} \rightarrow Y_0$$

Is the simple bound also valid for the small-mass range?

Abazajian, Fuller, Patel, 01'



## A Stationary State?

$$Y_{\nu_\tau} \rightarrow Y_0$$



$$\mathcal{I}_\nu \approx \mathcal{I}_{\bar{\nu}} \approx \mathcal{I}$$

Including the degen. param.


$$f_E^{\nu_\tau} = \frac{1}{\exp[E/T - \eta] + 1}$$

$$f_E^{\bar{\nu}_\tau} = \frac{1}{\exp[E/T + \eta] + 1}$$



# Energy-loss Rates and SN Bounds

## Criterion for a stationary state



## Evolution of the degeneracy parameter

$$\begin{aligned}
 \dot{N}_{\nu_\tau} &= -\frac{1}{4} \sum_a \int \frac{E^2 dE}{2\pi^2} \sin^2 2\vartheta_\nu \int \frac{E'^2 dE'}{2\pi^2} \mathcal{W}_{E'E}^a f_{E'}^{\nu_\tau} \\
 \dot{N}_{\bar{\nu}_\tau} &= -\frac{1}{4} \sum_a \int \frac{E^2 dE}{2\pi^2} \sin^2 2\vartheta_\nu \int \frac{E'^2 dE'}{2\pi^2} \overline{\mathcal{W}}_{E'E}^a f_{E'}^{\bar{\nu}_\tau}
 \end{aligned}$$

$$\begin{aligned}
 f_{E'}^{\nu_\tau} &= \frac{1}{\exp[E/T - \eta] + 1} \\
 f_{E'}^{\bar{\nu}_\tau} &= \frac{1}{\exp[E/T + \eta] + 1}
 \end{aligned}$$

Sterile neutrinos with mixing angles  $\vartheta_\nu < \vartheta_c \approx 10^2$  can escape from the core.

# Energy-loss Rates and SN Bounds

## Evolution of the degeneracy parameter

$$\frac{d}{dt}\eta(t) = \frac{N_B G_F^2 s_{2\theta}^2 T^2}{4\pi} [\mathcal{F}_{\bar{\nu}}(\eta) - \mathcal{F}_{\nu}(\eta)] \mathcal{G}^{-1}(\eta)$$

## Feedback effects

Initial condition:  $t = 0, \eta = 0$

$$\mathcal{F}_{\bar{\nu}}(0) - \mathcal{F}_{\nu}(0) > 0$$

Antineutrino Emission Rate  $>$  Neutrino Emission Rate

$Y_{\nu_\tau}$  ( $\tau$  number asymmetry)  $\uparrow$

$$E_r \propto |Y_0 - Y_{\nu_\tau}|^{-1} \uparrow$$

$\mathcal{G}_{\bar{\nu}}$   $\downarrow$

$\mathcal{G}_{\nu}$   $\uparrow$

$\eta$  increases ( $\eta > 0$ )

$$f_E^{\nu_\tau} = \frac{1}{\exp[E/T - \eta] + 1}$$

$$f_E^{\bar{\nu}_\tau} = \frac{1}{\exp[E/T + \eta] + 1}$$

Antineutrino Emission Rate  $\downarrow$

Neutrino Emission Rate  $\uparrow$

stable point  $\eta^*$

$$\mathcal{F}_{\bar{\nu}}(\eta^*) = \mathcal{F}_{\nu}(\eta^*)$$

Neutrino Emission Rate

=



Antineutrino Emission Rate

# Energy-loss Rates and SN Bounds

## Evolution of the degeneracy parameter



Raffelt, Zhou, 11'

$$\frac{d}{dt} \eta(t) = \frac{N_B G_F^2 s_{2g}^2 T^2}{4\pi} [\mathcal{F}_\nu(\eta) - \mathcal{F}_\nu(\eta)] \mathcal{G}^{-1}(\eta)$$



# Energy-loss Rates and SN Bounds

1. The stable point  $\eta^*$  can be either negative or positive, depending on the sterile neutrino mass and vacuum mixing angle;
2. The values of  $\eta^*$  are negative for large vacuum mixing angles, because more antineutrinos than neutrinos are trapped in the SN core;
3. We temporarily ignore the trapped sterile neutrinos, which may actually transfer energies rapidly due to their larger mean free paths.



# Energy-loss Rates and SN Bounds

Raffelt, Zhou, 11'

## Energy-loss rate

$$\mathcal{E}_s(t) = \frac{N_B G_F^2 s_{2g}^2 T^6}{8\pi^3} [\mathcal{R}_{\bar{\nu}}(\eta) + \mathcal{R}_{\nu}(\eta)]$$


$$\mathcal{R}_{\bar{\nu}}(\eta) = \int_0^\infty \frac{x^5}{e^{x+\eta} + 1} \frac{1 - \mathcal{B}(x, x_r \varepsilon^-, x_r \varepsilon^+)}{s_{2g}^2 + (c_{2g} - x/x_r)^2} dx$$

## Averaged energy-loss rate

$$\langle \mathcal{E}_s \rangle = \tau_d^{-1} \int_0^{\tau_d} \mathcal{E}_s(t) dt$$

## Supernova bound

$$\langle \mathcal{E}_s \rangle < 3.0 \times 10^{33} \text{ erg cm}^{-3} \text{ s}^{-1}$$




**Thank you for your attention!**



# Sterile Neutrinos and SN Explosions

## Mixing with electron neutrinos



## How to constrain sterile neutrinos?

### Remarks:

1. If the lepton-number loss is not significant, one can simply apply the standard energy-loss argument to the  $\nu_e$ - $\nu_s$  mixing case;
2. For the warm-dark-matter mass range (1 keV to 10 keV), the MSW resonance may be present and amplify the lepton-number-loss rate;
3. Sterile neutrinos have already done something important during the collapsing phase, such as reducing the electron number fraction  $Y_e$  and thus the size of the homologous core, and the energy of the shock wave.



# Sterile Neutrinos and SN Explosions

## Sterile neutrino assisted SN explosions?

Hidaka, Fuller, 06'

One-zone model of the collapsing core: the EoS & resonant  $\nu_e$ - $\nu_s$  conversion, ...

To include the neutrino trapping and diffusion, shock-wave propagation, ...

